

BURST NEUTRINOS FROM NITROGEN FLASH

A. M. SERENELLI

Institute for Advanced Study, Einstein Drive, Princeton, NJ 08540, USA

AND

M. FUKUGITA

Institute for Advanced Study, Einstein Drive, Princeton, NJ 08540, USA and
 Institute for Cosmic Ray Research, University of Tokyo, Kashiwa 277-8582, Japan

Draft version February 2, 2008

ABSTRACT

Neutrinos give a novel probe to explore deep interior of astrophysical objects, which otherwise is not accessible with optical observations; among notable examples are solar and supernova neutrinos. We show that there is a new class of strong neutrino emission from helium burning, $^{14}\text{N} + \alpha \rightarrow ^{18}\text{F} + \gamma$ followed by beta decay $^{18}\text{F} \rightarrow ^{18}\text{O} + e^+ + \nu_e$, that gives a maximum neutrino luminosity of 10^8 times the solar bolometric luminosity at the helium-core flash of a $1 M_\odot$ star, whereas the flash is not observable by optical means. This means that the neutrino flux, of average energy of 0.382 MeV, will be 10% the solar CNO neutrino flux on Earth if the star is located at 10pc.

Subject headings: stars: evolution — neutrinos

1. PRELIMINARIES

Low mass stars ($M < 2M_\odot$) are known to develop a degenerate helium core during their evolution along the Red Giant Branch (see for instance Sweigart & Gross 1978). The core is fed by the surrounding hydrogen-burning shell that slowly moves outwards in mass as hydrogen burning proceeds through the CNO bi-cycle. The helium mass fraction in the hydrogen-depleted core is $Y = 1 - Z$, where Z is the total heavy element mass fraction (note that Z does not change during hydrogen burning). As a result of the CNO burning, Z is composed by approximately 70% of ^{14}N . When the mass of the degenerate core reaches about $0.47M_\odot$, the temperature rises up to 10^8K and helium ignites under degenerate conditions. At the moment of helium ignition, the total number of ^{14}N nuclei in the core is $n(^{14}\text{N}) \approx 5.3 \times 10^{53} (M_c/0.47) (Z/0.019)$, where M_c is the core mass in solar mass units. At this temperature destruction of ^{14}N during helium burning (independently of the stellar mass) occurs through the reaction $^{14}\text{N} + \alpha \rightarrow ^{18}\text{F} + \gamma$ followed by beta decay $^{18}\text{F} \rightarrow ^{18}\text{O} + e^+ + \nu_e$. The maximum energy of these neutrinos is 0.633 MeV, while the average energy 0.382 MeV. This implies that were all the ^{14}N burned as described above, a total energy $\epsilon_\nu \approx 3.2 \times 10^{47} (M_c/0.47) (Z/0.019)$ erg would be carried away from the star by neutrinos. The importance of this neutrino emission contrasts with the folklore that neutrino emission from nuclear reactions is not important during the helium burning phase. It is the purpose of this work to show, using stellar evolutionary calculations, that most of this energy is actually released in the very short time-scale that characterizes the helium-core flash and that the neutrino flux reaches values as high as $6 \times 10^{47} \text{ s}^{-1}$.

2. THE HELIUM-CORE FLASH

The detailed structure of the degenerate cores of low mass RGB stars at the moment of helium ignition has been widely discussed in the literature (Sweigart & Gross 1978; Sweigart et al. 1989; Salaris et al. 2002). Although

uncertainties in the modeling exist, a very good understanding of RGB stars has been achieved. This is reflected, for instance, in the work by Raffelt (1990) where it has been shown that observations of globular cluster stars and RR Lyrae stars indicate that the core mass at helium-flash ignition inferred from observations and those predicted by stellar evolution calculations agree at about a 5% level. This has been used to constrain different properties of neutrinos and also on some more exotic particles like axions (Raffelt & Weiss 1992, 1995).

In this work, however, we are particularly interested in nuclear neutrino production during the helium-core flash, which is a less well understood evolutionary phase. In particular, the central question that remains to be answered in a satisfactory way is whether the helium-core flash is a dynamic event and whether dynamical episodes are of any consequence to the flash itself and the later evolution. However, hydrodynamic 2-D and 3-D calculations (Deupree 1996; Achatz 1995; and Achatz, Müller & Weiss, in preparation) show that hydrostatic stellar evolution calculations appear to follow the global properties of the helium-core flash reasonably well, at least in its global properties. With the confidence gained from those results, we use for the present work the LPCODE stellar evolution code (Althaus et al. 2003). Here we review some important aspects of particular relevance for the present work. The nuclear network has been updated and now includes 33 isotopes from H to ^{32}P and 99 reactions (charged particles reaction rates are taken from Angulo et al. 1999). Mixing in convective regions is treated as a diffusive process and equations are fully coupled to nuclear burning. Conductive opacities are taken from Hubbard & Lampard (1969). Neutrino emission rates are from Itoh and collaborators, exception made of the plasma neutrino emission rate that has been adopted from Haft et al. (1994). More details can be found in Althaus et al. (2003).

In what follows, we present the results for the evolution of a $1 M_\odot$ stellar model of initial solar composition ($Y, Z = (0.275, 0.019)$), our reference calculation.

Let us first briefly discuss the evolution of the model up to the helium-core flash. Calculations were started from a fully convective pre-main sequence model. The main sequence phase lasts about 9.5 Gyr and when hydrogen is depleted in the core the star evolves to the RGB phase, where the degenerate helium-core grows as burning in the hydrogen shell proceeds. As the mass of the helium core increases, it contracts and releases gravitational energy that heats it up. Additionally, the temperature of the burning shell is mainly determined by the local value of the gravitational potential and thus increases steadily, also contributing to increase the core temperature. The dominating cooling mechanism in the core is the plasma neutrino emission, that also increases along with the core size. The net result is that the core temperature increases steadily along the RGB evolution, but neutrino emission is enhanced closer to the center in the almost isothermal core, eventually leading to a temperature inversion. In our evolutionary sequence, this happens when the degenerate core has $0.27M_{\odot}$ and the age and luminosity of the model are 12.62 Gyr and $\log(L/L_{\odot})=1.8$ respectively. The model continues to evolve along the RGB with its luminosity and core temperature increasing continuously. The core degeneracy also continues to increase, decoupling the pressure from the temperature. This continues until the temperature in the core is finally enough to ignite the helium and the evolution along the RGB is halted. This defines the tip of the RGB. In our calculations, the luminosity at the tip of the RGB for this model is $\log(L_{\text{tip}}/L_{\odot})=3.446$ and the degenerate helium-core has $0.476 M_{\odot}$. The age of the model is 12.664 Gyr. The temperature inversion is located at $M_{\text{max}} = 0.23M_{\odot}$; this sets the position where helium ignites. At the moment when the helium-burning luminosity L_{He} exceeds for the first time that of the hydrogen-burning, the temperature T_{max} and density ρ_{max} at M_{max} are 1.1×10^8 K and $3.2 \times 10^5 \text{ g cm}^{-3}$ respectively; the degeneracy parameter (Kippenhahn & Weigert 1990, Sect. 15.4) is $\psi = 8.5$.

Given the conditions at the moment of helium ignition, the core is supported by electronic degeneracy pressure and, as mentioned before, temperature is decoupled from pressure. The helium flash proceeds through the triple alpha (3α) reaction, that has a Maxwellian averaged rate $N_A^2 \langle \sigma v \rangle_{3\alpha}$ very strongly dependent on temperature: $N_A^2 \langle \sigma v \rangle_{3\alpha} \propto T^{40, \dots, 20}$ for $T = 1, \dots, 2.5 \times 10^8$ K. This implies that a small increase in temperature results in an enormous increase in the burning rate, and because the structure of the star cannot respond to an increase in T by expanding and cooling, a thermonuclear runaway, the helium-core flash, quickly follows helium ignition. Time evolution during the helium-core flash of some important quantities is shown in Fig. 1. The top panel presents the time evolution of helium burning luminosity L_{He} . In just a few years, L_{He} grows from being a negligible energy source in the star to overcome the luminosity of the star (L_{bol}) by almost seven orders of magnitude. It is important to note that when L_{He} reaches about $10^2 L_{\odot}$, energy cannot be transported by radiation alone, and adiabatic convection sets in in the layers above M_{max} . At its maximum extension, the convective shell ranges from M_{max} up to $0.475M_{\odot}$. As shown in the next section, the extension of this convective shell determines to a large extent the total amount of N^{14} that is actually

burnt during the flash. Fig. 1 also shows that the helium flash does not show up in L_{bol} , which actually decreases as the hydrogen burning luminosity (L_{H}) drops abruptly when the layers surrounding the core expand and cool. To illustrate the conditions under which the flash starts and evolves, the middle and lower panels in Fig. 1 show the temperature and density at M_{max} . The lower panel, in particular, makes evident that the energy release during the flash effectively removes the degeneracy in the burning region and is used to lift up the stellar core from its potential well.

After the main episode of the helium-core flash, described above, evolutionary calculations show that a series of sub-flashes follow, each of them occurring successively closer to the center until degeneracy is completely removed from the core and steady helium-burning is established Serenelli & Weiss (2005). Detailed evolution through this phase is rather uncertain, in particular because it is not well understood how heat flows from the point of maximum temperature inwards (Achatz 1995; Deupree 1996). However, we do not expect that this affects our main results, because the bulk of the nuclear neutrinos and in particular the sharp peak in the neutrino luminosity occurs during the main flash episode. We show this in the next section.

3. NEUTRINOS FROM THE NITROGEN FLASH

We now focus on the production of nuclear neutrinos during the helium-core flash. As mentioned before, ^{14}N is, by far, the second main constituent of the helium core at the time of helium flash ignition. The Maxwellian averaged rate of the $^{14}\text{N} + \alpha \rightarrow ^{18}\text{F} + \gamma$ reaction, $N_A \langle \sigma v \rangle_{14}$ (N_A is the Avogadro number), is between 3 and 5 orders of magnitude larger than that of the 3α reaction in the temperature range characteristic of the helium flash (Angulo et al. 1999) and its dependence on temperature is extremely high ($N_A \langle \sigma v \rangle_{14} \propto T^{45, \dots, 21}$ for $T = 1, \dots, 2.5 \times 10^8 \text{ K}$). The linear dependence on ρ of the $^{14}\text{N} + \alpha$ reaction delays its activation with respect to helium ignition through 3α , but the extreme temperature dependence leads to a very steep increase in the rate as temperature rises¹. The ^{14}N half-life for α -captures decreases from 10^{15} s when helium ignites at $T = 10^8 \text{ K}$ to 1.7×10^4 s at $T = 2.25 \times 10^8 \text{ K}$, the maximum temperature reached in the flash. We call this the “Nitrogen Flash”, a term coined by Iben (1967) in a pioneering work on helium-flash calculations. At that time, the $^{14}\text{N} + \alpha$ reaction was thought to halt the evolution along the Red Giant Branch and actually lead to the ignition of the helium-flash. It seems clear now that, as described above, this is not the case and ^{14}N burning lags slightly behind the development of the flash, but we believe the name is still appropriate.

The complete chain for ^{14}N burning is $^{14}\text{N} + \alpha \rightarrow ^{18}\text{F} + \gamma$ followed by beta decay $^{18}\text{F} \rightarrow ^{18}\text{O} + e^+ + \nu_e$, where the neutrinos are produced with a maximum energy of 0.633 MeV. The beta

¹ The helium flash is, at all times, driven by the 3α reaction, while ^{14}N burning supplies at most 10% of the nuclear energy release and for a very short time. This is so due to a combination of factors: when the flash starts $\rho > 10^5 \text{ g cm}^{-3}$ and 3α dominates because its rate depends as ρ^2 . At later times, when the flash develops and starts producing an expansion of the burning region, density drops but ^{14}N is rapidly depleted.

decay half-life of ^{18}F is only 6.6×10^3 s, shorter than the half-life of ^{14}N for α -captures. Other destruction channels of ^{18}F are at most two orders of magnitude slower than the beta decay and do not play any significant role in the ^{18}F nucleosynthesis (in particular $^{18}\text{F}(n, \alpha)^{15}\text{N}$ dominates among these channels when a small mass fraction of neutrons of about 10^{-19} builds up transiently at the base of the burning shell). This has the consequence that the neutrino production rate basically follows that of ^{14}N destruction. In the upper panel of Fig. 2 we show the time evolution of the nuclear neutrino luminosity produced by ^{18}F decay. Note the very short time-scale that dominates its evolution, more than 4 orders of magnitude increase in only 0.2 years. The maximum luminosity is about $10^8 L_\odot = 3.8 \times 10^{41} \text{ erg s}^{-1}$ and the resulting number flux of neutrinos at the peak is about $6 \times 10^{47} \text{ s}^{-1}$. We have shown five epochs of the internal distribution of the ^{14}N mass fraction in the lower panel of Fig. 2 in order to illustrate how ^{14}N depletion and, accordingly, neutrino production proceeds. ^{14}N in the shell of $M = 0.23$ to $0.475 M_\odot$ is destroyed nearly entirely within 1 yr. Indeed, $2.65 \times 10^{-3} M_\odot$ of ^{14}N , which is 50% of ^{14}N produced in the previous CNO burning is depleted in this short time-scale. The nuclear neutrino energy emitted in nitrogen flash amounts to 0.05% the value emitted in the entire life of the star. ^{18}O produced in the beta decay is eventually all consumed to produce ^{22}Ne by further α -captures, which has a reaction rate comparable to the $^{14}\text{N} + \alpha$ reaction, and therefore it entirely disappears in the core of white dwarfs. The only in-principle visible event that signatures helium flash are the burst neutrinos.

The peak luminosity of nuclear neutrinos during the helium-core flash (L_ν^{peak}) is directly linked to the strength of the helium-flash and thus uncertainties in the helium-flash calculations will also affect the value of (L_ν^{peak}). A number of factors contribute to the uncertainty in the helium flash calculations. On one hand, uncertainties in the input physics is amply dominated by uncertainties of a factor of 2 in the electron conductive opacities (κ_{cond}) in degenerate cores of RGB stars (Salaris et al. 2002; Catelan 2005).

Other possible sources of uncertainty like thermal neutrino emission, equation of state or radiative opacities are much better constrained and have a much smaller impact on the uncertainties of RGB stellar models (Salaris et al. 2002); and consequently in the nuclear neutrino emission. A second source of uncertainty, that cannot currently be quantified, is that introduced by the lack of complete hydrodynamic calculations of the helium-core flash. Although these calculations seems to support hydrodynamic stellar evolutionary calculations (in particular Achatz 1995 shows that no mixing beyond the canonical convective boundaries as given by the Schwarzschild criterion occur and also that the temperature gradient in the convective shell is very close to the adiabatic one), uncertainties in how heat flows below the temperature inversion are present and can have some influence on the temporal evolution of the temperature at M_{max} . Finally, the peak luminosity of nuclear neutrinos depends on the mass and composition of the star. We have addressed the first and last points by performing a series of additional evolutionary calculations. We do not attempt

to cover the parameter space, but just to have a rough estimation of the level of the uncertainties that can be expected. The results are summarized in Table 1. The behavior of L_ν^{peak} at peak on mass and helium content reflects that of L_{He} at peak. The behavior on the metallicity follows approximately the linear relation expected from the simple arguments presented in Sect. 1. The dependence of L_ν^{peak} on κ_{cond} seems to be linear, and a factor of 2 uncertainty in κ_{cond} translates into a factor of 2 uncertainty in L_ν^{peak} at peak. A last source of uncertainty, affecting only the neutrino production, is the $\text{N}^{14} + \alpha$ reaction rate. Its uncertainty directly translates into an uncertainty in L_ν^{peak} . Görres et al. (2000) have determined this reaction rate to be 2 to 3 times smaller than that of Angulo et al. 1999 (the value adopted in this work), with an uncertainty of about 20% in the temperature range of interest.

4. CONCLUDING REMARKS

We have shown that during the main episode of the helium-core flash a burst of neutrinos is produced due to the extremely rapid burning of N^{14} by α captures in the so-called Nitrogen Flash. Our reference calculation, a $1 M_\odot$ stellar model of solar composition (Y, Z) = (0.275, 0.019) gives a nuclear neutrino peak luminosity of about $10^8 L_\odot$. Neutrinos from the Nitrogen Flash carry precious information about the degenerate conditions under which the helium-core flash is believed to occur. They may be the only direct manifestation we can ever detect of such an event, that does not manifest itself in the stellar surface. The neutrino flux expected at the flash is $5.3 \times 10^7 (d/10\text{pc})^{-2} \text{ cm}^{-2}\text{s}^{-1}$ at maximum (for a star located d pc away), which lasts for 3 days. This is about 10% of the CNO neutrino flux from the Sun.

We are not optimistic about the detection of these burst neutrinos, not only because this is a rare event, but also because solar neutrinos will give a serious background. A very large volume detector with the capability of energy resolution with the threshold set just above the maximum energy of p-p neutrinos is needed. To give an idea, in a 1000 ton hydrocarbon scintillator, we expect for a helium flash at 10 pc 10 events for 3 days, which is to be compared to 80 events from solar CNO neutrinos for the neutrino energy window of 0.42 to 0.63 Mev (here neutrino oscillation is ignored). In practice, we also expect a background from solar ^7Be neutrinos for electron scattering.

We want to acknowledge James Gunn and Bohdan Paczyński for valuable discussions and comments, John Bahcall and Achim Weiss for a careful reading of the manuscript, and Alan Chen for pointing out the latest reference on the $\text{N}^{14} + \alpha$ reaction rate. We also welcome comments from the referee, that have helped to improve the original manuscript substantially. A.M.S. is supported in part by the NSF through the grant PHY-0070928 to the Institute for Advanced Study and M.F. in part by Monell Foundation at Princeton.

REFERENCES

- Achatz, K. 1995, Master's Thesis, Tech. Univ. Munich
 Angulo, C. et al. 1999, Nuc. Phys. A, 656, 3
 Althaus, L. G., Serenelli, A. M., Córscico, A. H., & Montgomery, M. H. 2003, A&A, 404, 593
 Catelan, M. 2005, in ASP Conf. Ser. TBA, Resolved Stellar Populations, ed. D. Valls-Gabaud, & M. Chavez, to be published, preprint (astro-ph/0507464)
 Deupree, R. G. 1996, ApJ, 471, 377
 Görres, J., Arlandini, C., Giesen, U., Heil, M., Kppeler, F., Leiste, H., Stech, E., & Wiescher, M. 2000, Phys. Rev. C, 62, 055801
 Haft, M., Raffelt, G., & Weiss, A. 1994, ApJ, 425, 222
 Hubbard, W. B., & Lampe, M. 1969, ApJS, 19, 297
 Iben, I., Jr. 1967, ApJ, 147, 650
 Itoh, N., Mitake, S., Iyetomi, H., & Ichimaru, S. 1983, ApJ, 273, 774
 Kippenhahn, R. & Weigert A. 1990, in *Stellar Structure and Evolution*, Springer-Verlag, Berlin
 Raffelt, G. G. 1990, ApJ, 365, 559
 Raffelt, G. G., & Weiss, A. 1992, A&A, 264, 536
 Raffelt, G. G., & Weiss, A. 1995, Phys. Rev. D, 51, 1495
 Salaris, M., Cassisi, S., & Weiss, A. 2002, PASP, 114, 375
 Schwarzschild, M., & Härm, R. 1962, ApJ, 136, 158
 Serenelli, A. M., & Weiss, A. 2005, A&A, accepted for publication, preprint (astro-ph/0507430)
 Sweigart, A. V., Greggio, L., & Renzini, A. 1989, ApJS, 69, 911
 Sweigart, A. V., & Gross, P. G. 1978, ApJ, 36, 405

TABLE 1

IMPORTANT QUANTITIES CHARACTERIZING THE HELIUM-CORE FLASH OF OUR EVOLUTIONARY SEQUENCES.

Model	M_c	$\log L_{\text{Tip}}$	$\log L_{\text{He}}$	$\log L_\nu$
Reference	0.476	3.446	9.84	7.95
$0.5 \times \kappa_{\text{cond}}$	0.484	3.494	10.18	8.29
$2 \times \kappa_{\text{cond}}$	0.468	3.403	9.58	7.65
$Y = 0.2505$	0.480	3.440	9.97	8.12
$Y = 0.3005$	0.472	3.430	9.71	7.78
$Z = 0.029$	0.474	3.442	9.78	8.07
$Z = 0.009$	0.479	3.413	9.90	7.70
$Z = 0.0019$	0.482	3.347	10.02	7.14
$M = 0.80$	0.478	3.432	9.88	8.01
$M = 1.20$	0.476	3.434	9.83	7.94
$M = 1.40$	0.476	3.434	9.86	7.95

NOTE. — The table summarizes the most relevant quantities of our evolutionary sequences. First column denotes the model by giving the parameter that has been modified with respect to the reference model (first row). Second and third columns give the helium-core size and luminosity at the RGB-tip. Fourth and fifth columns give the peak helium-burning and nuclear neutrino luminosities during the flash. In all cases luminosities are in solar units.




FIG. 1.— Temporal evolution of stellar quantities during the development of the helium-core flash in our $1 M_\odot$ stellar model. Top panel shows the evolution of helium (L_{He}) and hydrogen (L_{H}) burning luminosities, and stellar luminosity (L_{bol}) in solar units ($L_\odot = 3.84 \times 10^{33} \text{ erg s}^{-1}$). The stellar luminosity has been shifted up by 0.2 dex to make the otherwise two degenerate lines (L_{H} and L_{bol}) visible. Middle and lower panels show, respectively, the temperature and density at the point of maximum temperature in the models (M_{max}), located off-center, at about $0.23 M_\odot$. $\tau = 0$ corresponds to the moment of maximum L_{He} .




FIG. 2.— Top panel. The evolution of the nuclear neutrinos luminosity (L_ν) very close to the flash peak ($\tau = 0$). Note that nuclear neutrinos during the evolutionary phase shown here are solely created in the beta decay $^{18}\text{F} \rightarrow e^+ + \nu_e$ that follows the α -capture $^{14}\text{N} + \alpha \rightarrow ^{18}\text{F} + \gamma$. Vertical lines, labeled A, B, C, D and E denote five chosen epochs for which the internal abundance profile of ^{14}N in mass fraction is shown in the lower panel. Initially (A) ^{14}N is barely depleted. The temperature dependence of the reaction $^{14}\text{N}(\alpha, \gamma)^{18}\text{F}$ is even larger than that of the 3α reaction and when $^{14}\text{N}(\alpha, \gamma)^{18}\text{F}$ becomes fully active it produces a burst of neutrinos that lasts for less than 4 days (B-C) during which the neutrino luminosity of the star reaches a maximum value of about $10^8 L_\odot$ (this is about a 5% percent of the total luminosity of nuclear neutrinos of the Milky Way). 50% of the available ^{14}N in the burning region is burnt in this time interval. Strong ^{14}N burning continues (D) but removal of degeneracy and subsequent expansion and cooling of the burning layers quickly follow and set a much slower pace for ^{14}N depletion (E). The step-like shape of the profiles denotes the presence of convection, that keeps the shell with an almost homogeneous composition even when nuclear burning is actually occurring only in the $0.23 - 0.27 M_\odot$ region. In passing by, we note that the shape of the ^{14}N profile in the inner $0.2 M_\odot$ reflects the increasing temperature of hydrogen-shell burning with increasing core size during previous evolutionary stages.